

NUTRISON 800 COMPLETE MULTI FIBRE

A nutritionally complete, high protein, lower energy, fibre enriched, ready-to-use enteral tube feed.

FEATURES

- **Suitable as a sole source of nutrition[^]**
- **0.83kcal/ml:** for patients requiring lower energy diets.¹⁻³
- **55g protein (27%E) per 1000ml:** to meet the protein requirements of patients with low energy requirements.
- **15g of MF6* fibre blend per 1000ml:** to help maintain normal bowel function.
- **Fish oils:** providing Docosahexaenoic acid (DHA) and Eicosapentaenoic acid (EPA).
- **Medium chain triglycerides (MCT):** for easier fat digestion and absorption.⁴⁻⁵
- **Whey-dominant P4 protein blend:** in line with international recommendations on protein quality/ amino acid profile and for gastro-intestinal tolerance benefits.⁶⁻¹³
- **Enriched with carotenoids:** in line with general health recommendations for their antioxidant properties and positive effect on immune function.¹⁴
- **1000ml OpTri bottle:** suitable for closed or open system feeding via ISO compliant flip-top screw cap.

Indications

For the dietary management of:

- Disease related malnutrition.
- Patients with reduced energy requirements (e.g. patients who are immobile and/or have a severe neurological disorder).

Important Notice

- Not for parenteral use.
- Not suitable for patients requiring a fibre free diet.
- Not suitable for patients with galactosaemia.
- Not suitable for patients with cow's milk protein allergy.
- Not suitable for infants under 1 year of age
- Use with caution in children aged 1-12 years of age.
- Use with caution in individuals with a seafood allergy.

Directions for Use

- Shake well before use.
- Use at room temperature.
- Handle aseptically to ensure product remains sterile.
- Usage to be determined by a healthcare professional.

Storage

- Store in a cool, dry place.
- Once opened, store in the refrigerator.
- Discard unused contents after 24 hours.

^{*}MF6 is a unique, patented blend of six soluble and insoluble fibres (soy polysaccharide, cellulose, resistant starch, gum arabic, oligofructose and inulin) reflecting the proportions of the different fibre types in a healthy diet.

Ordering Information

To order contact Nutricia Customer Care **0800 688 747**.


Nutrison 800 Complete Multi Fibre	Product code	Units per carton	Pharmacode
1000ml OpTri bottle	132379	8	2572982

Ingredients

Nutrison 800 Complete Multi Fibre: Water, maltodextrin, whey protein (from **milk**), vegetable oils (sunflower oil, rapeseed oil, MCT oil (coconut oil, palm kernel oil)), dietary fibres (inulin, oligofructose, arabic gum, **soy** polysaccharides, cellulose, resistant starch), **milk** protein caseinate, pea protein, **soy** protein, rice flour, potassium citrate, sodium chloride, magnesium hydrogen phosphate, calcium carbonate, **fish oil**, emulsifier (**soy** lecithin), tri calcium phosphate, potassium hydroxide, carotenoids (contain **soy**) (β -carotene, lutein, lycopene oleoresin from tomatoes), choline chloride, sodium L-ascorbate, tri sodium citrate, ferrous lactate, zinc sulphate, nicotinamide, DL- α tocopheryl acetate, retinyl acetate, copper gluconate, manganese sulphate, sodium selenite, calcium D-pantothenate, chromium chloride, D-biotin, cholecalciferol, thiamin hydrochloride, pteroylmonoglutamic acid, pyridoxine hydrochloride, riboflavin, potassium iodide, sodium fluoride, sodium molybdate, phytomenadione, cyanocobalamin.

Allergen & Cultural Information

- Contains: cow's milk protein, soy, fish oil.
- Does not contain: wheat, egg, nuts,** lupins.
- Halal certified.
- No Kosher forbidden ingredients.
- No gluten containing ingredients. No detectable gluten when tested to a sensitivity level of less than 5 parts per million (<5 ppm i.e. <5mg/kg).
- Low lactose (lactose <2g/100g).



NUTRISON 800 COMPLETE MULTI FIBRE

NUTRITION INFORMATION		Per 100ml	Per 1000ml
Energy	kcal	83	830
	kJ	345	3450
Protein	g	5.5 (27% E)	55
Casein	g	1.4	14
Whey	g	1.9	19
Soy	g	1.1	11
Pea	g	1.1	11
Carbohydrate	g	8.8 (43% E)	88
Sugars	g	0.6	6
as Lactose	g	<0.025	<0.25
Fat	g	2.5 (27%E)	25
Saturates	g	0.7	7
- of which MCT	g	0.4	4
Monounsaturates	g	1.3	13
Polyunsaturates	g	0.5	5
DHA	mg	20.3	203
EPA	mg	30.0	300
ω6 / ω3 ratio		2.3:1	2.3:1
Fibre	g	1.5 (3%E)	15
Soluble:Insoluble		80:20	80:20
Water	ml	86	860
Minerals		Per 100ml	Per 1000ml
Sodium	mg	150	1500
	mmol	6.5	65
Potassium	mg	225	2250
	mmol	5.8	58
Calcium	mg	120	1200
Phosphorus	mg	96	960
Magnesium	mg	27	270
Chloride	mg	145	1450
Ca:P ratio		1.2:1	1.2:1

Vitamins		Per 100ml	Per 1000ml
Vitamin A	µg-RE	123	1230
Vitamin D	µg	2.0	20
Vitamin E	mg α-TE	1.9	19
Vitamin K	µg	8.0	80
Vitamin C	mg	15	150
Thiamin	mg	0.23	2.3
Riboflavin	mg	0.24	2.4
Niacin	mg NE	2.7	27
Vitamin B6	mg	0.26	2.6
Vitamin B12	µg	0.32	3.2
Folic Acid	µg	40	400
Pantothenic Acid	mg	0.8	8
Biotin	µg	6.0	60
Trace Elements		Per 100ml	Per 1000ml
Iron	mg	2.4	24
Zinc	mg	1.8	18
Manganese	mg	0.5	5
Copper	µg	270	2700
Iodine	µg	20	200
Molybdenum	µg	15	150
Selenium	µg	8.5	85
Chromium	µg	10	100
Fluoride	mg	0.15	1.5
Other		Per 100ml	Per 1000ml
Carotenoids	mg	0.3	3
Choline	mg	55	550
Osmolality	mOsmol/ kgH ₂ O	325	325

[^] In accordance with Australia New Zealand Food Standards Code – Standard 2.9.5

^{**} Peanut (*Arachis hypogaea*), Almond (*Amygdalus communis* L. J.), Hazelnut (*Corylus avellana*), Walnut (*Juglans regia*), Cashew (*Anacardium occidentale*), Pecan nut (*Carya illinoensis* (Wangenh.) K. Koch), Brazil nut (*Bertholletia excelsa*), Pistachio nut (*Pistacia vera*), Macadamia nut and Queensland nut (*Macadamia ternifolia*) and products thereof.

REFERENCES 1. Dickerson RN *et al.* Measured energy expenditure of tube-fed patients with severe neurodevelopmental disabilities. *J Am Coll Nutr.* 1999 Feb;18(1):61-8. 2. Dickerson RN *et al.* Energy requirements of non-ambulatory, tube-fed adult patients with cerebral palsy and chronic hypothermia. *Nutrition.* 2003 Sep;19(9):741-6. 3. Dickerson RN *et al.* Validation of a new method for estimating resting energy expenditure of non-ambulatory tube-fed patients with severe neurodevelopmental disabilities. *Nutrition.* 2002 Jul-Aug;18(7-8):578-82. 4. Beckers EJ, Jeukendrup AE *et al.* Gastric emptying of carbohydrate--medium chain triglyceride suspensions at rest. *Int J Sports Med.* 1992 Nov;13(8):581-4. 5. Hunt JN, Knox MT. A relation between the chain length of fatty acids and the slowing of gastric emptying. *J Physiol.* 1968 Feb;194(2):327-36. 6. Hurt RT, McClave SA, Martindale RG, *et al.* Summary Points and Consensus Recommendations From the International Protein Summit. *Nutrition in Clinical Practice.* 2017;32:142S-151S. 7. World Health Organization. Protein and amino acid requirements in human nutrition: report of a joint FAO/WHO/UNU expert consultation. 2007; WHO technical report series ; no. 935. 8. Kuyumcu S, Menne D, Curcic J, *et al.* Noncoagulating enteral formula can empty faster from the stomach: A double-blind, randomized crossover trial using magnetic resonance imaging. *Journal of Parenteral and Enteral Nutrition.* 2015;39:544-551. 9. van den Braak CC, Klebach M, Abrahamse E, *et al.* A novel protein mixture containing vegetable proteins renders enteral nutrition products non-coagulating after in vitro gastric digestion. *Clinical Nutrition.* 2013;32:765-771. 10. Klebach M, Hofman Z, Bluemel S, *et al.* Effect of protein type in enteral nutrition formulas on coagulation in the stomach in vivo: Post hoc analyses of a randomized controlled trial with MRI. Abstract presented at Clinical Nutrition Week, January 16-19, Austin, Tx. *Journal of Parenteral and Enteral Nutrition.* 2016;40:134(21). 11. Lutikhoud J, van Norren K, Rijna H, *et al.* Jejunal feeding is followed by a greater rise in plasma cholecystokinin, peptide YY, glucagon-like peptide 1, and glucagon-like peptide 2 concentrations compared with gastric feeding in vivo in humans: a randomized trial. *Am J Clin Nutr.* 2016;103:435-43. 12. Abrahamse E, van der Lee S, van den Braak S, *et al.* Gastric non-coagulation of enteral tube feed yields faster gastric emptying of protein in a dynamic in vitro model. Abstract presented at 34th ESPEN Congress, Sept 8-11; Barcelona, Spain. *Clinical Nutrition Supplements.* 2012;7:PP239(119). 13. Liu J, Klebach M, Abrahamse E, *et al.* Specific protein mixture reduces coagulation: An in vitro stomach model study mimicking a gastric condition in critically ill patients. Poster presented at 38th ESPEN Congress, 17-20 September; Copenhagen, Denmark. *Clinical Nutrition.* 2016;35:MON-P182 (S220). 14. Cooper DA, Eldridge AL, Peters JC. Dietary carotenoids and certain cancers, heart disease and age-related macular degeneration: A review of recent research. *Nutrition Reviews.* 1999; 57: 201-214.

**A food for special medical purposes;
to be used under strict medical supervision.**

For more information call the
Nutricia Clinical Care Line 0800 636 228

NUTRICIA
LIFE-TRANSFORMING NUTRITION

Nutricia Ltd.
124 Favona Road,
Favona, Auckland, 2024, New Zealand.
www.nutriciamedical.co.nz
NC4697-Jun20 / NUT1165