

DIALAMINE

DESCRIPTION

A powder containing carbohydrate, essential amino acids, flavouring and low levels of minerals.

INDICATIONS

For the dietary management of advanced chronic renal failure and certain UCD, where an essential amino acid supplement is required.

PREPARATION AND ADMINISTRATION

Dialamine may be taken as a drink or a paste. Suggested dilution is 1 to 5, i.e. 20g of powder plus 100ml of water. Mix the prescribed amount of powder to a paste with cold water and make up to the required volume. If taken as a paste, water or diluted drinks should be offered at the same time.

ADMINISTRATION GUIDELINES

The dosage should be determined by a clinician or dietitian only and is dependent on the age, body weight and medical condition of the patient.

GENERAL PRECAUTIONS

Use under medical supervision. Not for parenteral use. Not suitable for use as a sole source of nutrition. Not suitable for infants less than six months old.

STORAGE

Store in a cool, dry place. Always replace the container lid after use. Once opened, use within one month. Once reconstituted, store in a refrigerator and consume within 24 hours. Stir or shake immediately before serving.

PACK SIZE

400g tin.

INGREDIENTS

Sugar, Dried Glucose Syrup, L-Lysine Acetate, Citric Acid, L-Leucine, L-Valine, L-Threonine, L-Isoleucine, L-Tyrosine, L-Phenylalanine, Flavouring, L-Cystine, L-Histidine, L-Methionine, L-Tryptophan, L-Ascorbic Acid, Colouring (E160(a), E162).

AVERAGE CONTENTS	UNIT	PER 100G	PER 100ML*	
Energy	kcal	360	62.1	
	kJ	1530	264	
Protein	g	25	4.3	
Carbohydrate	g	65	11.2	
	sugars	g	59	10.2
	lactose	g	-	-
Fat	g	nil added	nil added	
Dietary fibre	g	nil added	nil added	
Minerals				
sodium	mg (mmol)	< 10 (<0.4)	< 1.7 (<0.07)	
potassium	mg (mmol)	< 10 (<0.3)	< 1.7 (<0.05)	
chloride	mg (mmol)	< 20 (<0.6)	<3.5 (<0.1)	
calcium	mg (mmol)	< 5 (<0.1)	<0.9 (<0.02)	
phosphorus	mg (mmol)	< 5 (<0.2)	<0.9 (<0.03)	
magnesium	mg (mmol)	< 5 (<0.2)	<0.9 (<0.04)	
Vitamins				
vitamin C	mg	125	21.6	
Water				
osmolality	mOsm/kg H ₂ O	-	900	
Amino acid profile				
L-Cystine	g	1.2	0.2	
L-Histidine	g	1.2	0.2	
L-Isoleucine	g	3.3	0.57	
L-Leucine	g	5.13	0.89	
L-Lysine	g	4.2	0.72	
L-Methionine	g	1.2	0.2	
L-Phenylalanine	g	1.8	0.31	
L-Threonine	g	3.6	0.62	
L-Tryptophan	g	0.75	0.13	
L-Tyrosine	g	3	0.52	
L-Valine	g	4.62	0.8	

* 1 in 5 dilution



A food for special medical purposes; must be used under strict medical supervision.

For more information contact the Nutricia Care Line:
Australia: 1800 060 051
New Zealand: 0800 636 228
ncl@nutricia.com

NUTRICIA | METABOLICS
LIFE-TRANSFORMING NUTRITION